

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Not only priming: Soil microbiota may protect tomato from root pathogens

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1677702> since 2018-11-05T14:37:59Z

*Published version:*

DOI:10.1080/15592324.2018.1464855

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

Matteo Chialva, Yang Zhou, Davide Spadaro, Paola Bonfante

Not only priming: Soil microbiota may protect tomato from root pathogens

Plant Signaling & Behavior, VOL. 13, NO. 8, e1464855, 2018, (3 pages), 10.1080/15592324.2018.1464855

**The publisher's version is available at:**

<https://www.tandfonline.com/doi/full/10.1080/15592324.2018.1464855>

**When citing, please refer to the published version.**

**Link to this full text:**

<http://hdl.handle.net/2318/1677702>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

# Not Only Priming: Soil Microbiota May Protect Tomato from Root Pathogens

Matteo Chialva<sup>1</sup>, Yang Zhou<sup>1,2</sup>, Davide Spadaro<sup>3</sup> and Paola Bonfante<sup>1</sup>

<sup>1</sup>Department of Life Sciences and Systems Biology, University of Torino, Viale P.A. Mattioli 25, I-10125 Torino, Italy; <sup>2</sup>College of Horticulture, South China Agricultural University, No. 483, Wushan St., Tianhe Dist. Guangzhou, China PR, 510642; <sup>3</sup>Department of Agricultural, Forestry and Food Sciences (DISAFA) and AGROINNOVA, Centre of Competence for the Innovation in the Agroenvironmental Sector, University of Torino, Largo Braccini 2, I-10095 Grugliasco, Italy.

**Submitted:** 28 March 2018

**Accepted:**

**Keywords:** *Fusarium oxysporum* f. sp. *lycopersici*; arbuscular mycorrhizal fungi; defence responses; lignin biosynthesis; microbiota; suppressive and conducive soils; susceptible and resistant genotypes; tomato; gene expression.

Correspondence to:

Paola Bonfante

Email: [paola.bonfante@unito.it](mailto:paola.bonfante@unito.it)

## Abstract

An increasing number of studies have investigated soil microbial biodiversity. However, the mechanisms regulating plant responses to soil microbiota are largely unknown. A previous work tested the hypothesis that tomato plants grown on native soils with their complex microbiotas

respond differently from tomato growing in a sterile substrate. Two soils, suppressive or conducive to *Fusarium oxysporum* f. sp. *lycopersici* (FOL), and two genotypes susceptible and resistant to the same pathogen were considered. The work highlighted that the two tested soil microbiotas, irrespectively of their taxonomic composition, elicit the PAMP-triggered Immunity Pathway, the first level of plant defence, as well as an increased lignin synthesis, leading to an active protection when FOL is present in the soil. Here, we tested the expression of a panel of genes involved in Effector-Triggered Immunity (ETI), demonstrating that soil microbiota, beside genotype, affects plant resistance to FOL also modulating this pathway.

## TEXT

Next-generation sequencing (NGS) has enabled in-depth investigations of the microbial communities associated with animals, plants, and fungi. The awareness that multicellular eukaryotes host thousands of microbes, many beneficial, some essential and only a few deleterious has led to a paradigm shift in our knowledge of microbial–eukaryote interactions. NGS approaches helped us to reply to basic questions of traditional microbiology, as: ‘Which are the microbes thriving in that niche?’, and ‘What are they doing?’. Focusing on the plant side and starting from the pioneering researches by Bulgarelli et al.<sup>1</sup> and Lundberg et al.<sup>2</sup>, many other studies revealed the extraordinary diversity of microbes present on both roots, shoots, leaves, fruits<sup>3,4</sup>, and demonstrated how different parameters affect the composition of the microbiota: plant genotype, soil features, environmental parameters<sup>5,6</sup>. Interestingly, the environment resulted to be the driving force also for human microbiota, where it dominates over host genetics in shaping human gut microbiota<sup>7</sup>. The strict relationship existing between microbiota and their eukaryotic host has also led to the development of the *holobiont* concept<sup>8,9</sup>. Host-microbial systems, being a complex assembly of diverse organisms, constitute unique biological entities, defined as ‘meta-organisms’ or holobionts<sup>10</sup>. However, metagenomic sequencing has only given indirect responses to the questions opened by

these new scenarios: ‘How the host responds to its extended microbiota, which represents its second genome?’.

Chialva et al.<sup>11</sup> focused on tomato (*Solanum lycopersicum*), testing the hypothesis that plants grown on native soils display different responses to soil microbiotas. Using transcriptomics, proteomics, and biochemistry, the study has described the responses of two tomato genotypes (susceptible or resistant to FOL) grown on two native soils (conducive and suppressive to FOL) and an artificial substrate. Results showed that native soils, particularly the suppressive one, affect tomato responses by modulating pathways involved in responses to oxidative stress, phenol biosynthesis, lignin deposition, and PAMP-triggered Immunity (PTI). By contrast, in tomato plants grown on steam-disinfected soils, total phenols and PTI responses significantly decreased, suggesting a crucial role of soil microbiota in eliciting a priming effect. To validate those observations, the mycorrhizal fungus *Funnelliformis mosseae*, was selected as one of the most abundant AM fungi in both soils, and inoculated in tomato growing on steam-disinfected soils: the fungal inoculation partly rescued some of the local and systemic responses, which were identified as a part of the priming response.

Martinez-Medina et al.<sup>12</sup> have neatly identified different conditions where plant defence priming takes place and have acknowledged many beneficial microbes as a source for priming stimuli. Indeed, under the tested experimental conditions (native soils vs sterile substrate), tomato activates several genes involved in PTI, such as those encoding for PR proteins, WRKY transcription factors, ROS burst signalling and calcium signalling, which are involved in immune response<sup>13</sup>. To understand whether such an adaptive measure leads the plant to an enhanced defence readiness<sup>11</sup> tomato plants were inoculated with FOL. As expected, reduced disease symptoms were detected in the resistant genotype ('Battito') in both soils; but surprisingly the susceptible genotype 'Cuore di Bue' was partially protected from FOL on the suppressive soil. However, it is still unknown whether

the Effector-Triggered Immunity (ETI), *i.e.* the second barrier against pathogens, responds to soil microbiota.

Here, we hypothesized that the priming status raised in tomato by soil microbiota could elicit the expression of genes directly involved in ETI in the presence of FOL. With this aim, we selected a panel of genes involved in the ETI pathway (Table 1) and tested their expression by using RT-qPCR in FOL-inoculated plant roots according to the set-up and methods described in Chialva et al.<sup>11</sup>.

Results indicate that soil microbiota promoted the ETI response of plants after FOL infection (Fig. 1): while in RNA-seq experiment, where FOL was not present, ETI genes were not differentially expressed, in FOL-inoculated plants RT-qPCR experiment detected gene modulation<sup>11</sup>. Both genotypes significantly upregulated the expression of *RIN4* ( $p < 0.05$ ) in both native soils compared to the control substrate. This protein is a target of type III pili effector proteins (virulence factors) from bacterial pathogens and interacts with RPS2 and RPM1 R protein leading to hypersensitive response<sup>14,15</sup>. Moreover, we tested the expression of two previously described ETI-marker genes<sup>16</sup> and found that one of them coding for a UDP-glucosyltransferase family 1 protein (UDP) is upregulated in both soils ( $p < 0.05$ ) with the exception of the susceptible cultivar in the conducive soil. However, the other marker gene tested (UDP1) did not show differential expression across conditions. By contrast, the expression of the *I-2* R gene, directly involved in FOL race 2<sup>17</sup>, was upregulated only in the resistant genotype grown in the suppressive soil, while it remained consistent for the susceptible genotype in all the substrates. These results suggest a synergy between the genotype (presence of Resistance genes), the soil biological features, and – mechanistically – the ETI response. The 'Cuore di Bue' susceptible genotype has a more modulated response: FOL-suppressive soil with its microbiota activates the ETI response, while this action is not elicited in

the conducive soil. This well explains the modulation of *I-2* R gene: to be activated, plant defences require the suppressive soil microbiota acting on the resistant genotype, while the synergy between these two conditions is not satisfied in the susceptible genotype. The hypothesis may have an experimental validation by the presence of many bio-control *Fusaria* strains isolated in the Albenga soil<sup>18</sup>.

Our previous experiments demonstrated that soil microbiota leads to a priming ('state of alert') in tomato eliciting the PTI, which represents the first level of plant defence. When challenged by a pathogen, the alerted plant activates a new set of more specific genes related to the ETI, which is the second specific defence level (Fig. 2). This mechanism leads to a partial protection from the pathogen attack, even in the absence of specific resistance genes (as for the cultivar 'Cuore di Bue'). The modulation of the ETI-related genes indicates that native soil microbiota also affects plant response to FOL via ETI, in addition to the crucial role played by the genotype. In conclusion, the investigation of the mechanisms operating in plants in native soils and in the presence of complex soil microbiota has revealed new unexpected responses. It seems that - just like humans - the tomato plant living in non-sterile conditions can better activate its immunity defence via the interaction with its microbiota.

#### **Disclosure of potential conflicts of interest**

No potential conflicts of interest were disclosed.

#### **Acknowledgments**

The authors thank the members of the Mycoplant Consortium for the support in FOL infection experiments.

123  
1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Funding**

This research has received funding from the Mycoplant project (Root Microbiome for Plant Health: dissecting the role of soil fungi, D15E13000350003, Fondazione Compagnia di San Paolo Torino) and by the European Union’s Horizon 2020 research and innovation programme under grant agreement No 727929 (A novel and integrated approach to increase multiple and combined stress tolerance in plants using tomato as a model – TOMRES). M.C. was funded by a TOMRES fellowship.

**References**

1. Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, et al. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature 2012; 488:91–5.
2. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, Rio TG del, et al. Defining the core Arabidopsis thaliana root microbiome. Nature 2012; 488:86–90.
3. Bai Y, Müller DB, Srinivas G, Garrido-Oter R, Potthoff E, Rott M, Dombrowski N, Münch PC, Spaepen S, Remus-Emsermann M, et al. Functional overlap of the Arabidopsis leaf and root microbiota. Nature 2015; 528:364–9.
4. Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG. Plant compartment and biogeography affect microbiome composition in cultivated and native Agave species. New Phytol 2016; 209:798–811.



5. Zgadzaj R, Garrido-Oter R, Jensen DB, Koprivova A, Schulze-Lefert P, Radutoiu S. Root nodule symbiosis in *Lotus japonicus* drives the establishment of distinctive rhizosphere, root, and nodule bacterial communities. *Proc Natl Acad Sci* 2016; 113:E7996–8005.
6. Hamonts Kelly, Trivedi Pankaj, Garg Anshu, Janitz Caroline, Grinyer Jasmine, Holford Paul, Botha Frederik C., Anderson Ian C., Singh Brajesh K. Field study reveals core plant microbiota and relative importance of their drivers. *Environ Microbiol* 2018; 20:124–40.
7. Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, Costea PI, Godneva A, Kalka IN, Bar N, et al. Environment dominates over host genetics in shaping human gut microbiota. *Nature* 2018; 555:210–5.
8. Bordenstein SR, Theis KR. Host Biology in Light of the Microbiome: Ten Principles of Holobionts and Hologenomes. *PLOS Biol* 2015; 13:e1002226.
9. Theis KR, Dheilly NM, Klassen JL, Brucker RM, Baines JF, Bosch TCG, Cryan JF, Gilbert SF, Goodnight CJ, Lloyd EA, et al. Getting the Hologenome Concept Right: an Eco-Evolutionary Framework for Hosts and Their Microbiomes. *mSystems* 2016; 1:e00028-16.
10. Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. The role of microorganisms in coral health, disease and evolution. *Nat Rev Microbiol* 2007; 5:355–62.
11. Chialva M, Salvioli di Fossalunga A, Daghino S, Ghignone S, Bagnaresi P, Chiapello M, Novero M, Spadaro D, Perotto S, Bonfante P. Native soils with their microbiotas elicit a state of alert in tomato plants. *New Phytol* 2018;
12. Martinez-Medina A, Flors V, Heil M, Mauch-Mani B, Pieterse CMJ, Pozo MJ, Ton J, van Dam NM, Conrath U. Recognizing Plant Defense Priming. *Trends Plant Sci* 2016; 21:818–22.
13. Chiang Y-H, Coaker G. Effector Triggered Immunity: NLR Immune Perception and Downstream Defense Responses. *Arab Book* 2015; 13:e0183.

14. Mackey D, Belkhadir Y, Alonso JM, Ecker JR, Dangl JL. Arabidopsis RIN4 is a target of the  
type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* 2003;  
112:379–89.
15. Liu J, Elmore JM, Coaker G. Investigating the functions of the RIN4 protein complex during  
plant innate immune responses. *Plant Signal Behav* 2009; 4:1107–10.
16. Pombo MA, Zheng Y, Fernandez-Pozo N, Dunham DM, Fei Z, Martin GB. Transcriptomic  
analysis reveals tomato genes whose expression is induced specifically during effector-  
triggered immunity and identifies the Epk1 protein kinase which is required for the host  
response to three bacterial effector proteins. *Genome Biol* 2014; 15. Available from:  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4223163/>
17. Simons G, Groenendijk J, Wijbrandi J, Reijans M, Groenen J, Diergaarde P, Van der Lee T,  
Bleeker M, Onstenk J, de Both M, et al. Dissection of the fusarium I2 gene cluster in tomato  
reveals six homologs and one active gene copy. *Plant Cell* 1998; 10:1055–68.
18. Poli A, Lazzari A, Prigione V, Voyron S, Spadaro D, Varese GC. Influence of plant genotype  
on the cultivable fungi associated to tomato rhizosphere and roots in different soils. *Fungal Biol*  
2016; 120:862–72.

141  
1  
142  
3  
4  
143  
5  
6  
144  
7  
8  
9  
145  
10  
11  
146  
12  
13  
14  
147  
15  
16  
148  
17  
18  
149  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure Legends**

**Figure 1.**

**RT-qPCR relative expression levels of gene involved in ETI in tomato plants (*Solanum lycopersicum*) infected with *Fusarium oxysporum* f. sp. *lycopersici* (FOL).**

*Ubiquitin* gene was used as reference for RT-qPCR. Letters indicate statistically supported differences (Kruskal–Wallis test at  $P < 0.05$ ). Data are means  $\pm$  SE ( $n = 3$ ). AL, ‘Albenga’ suppressive soil; RO, ‘Rosta’ conducive soil; CONT, Control 'Neutral' soil. B, 'Battito' FOL-resistant genotype; C, 'Cuore di Bue' FOL-susceptible genotype. (A) *RIN4*, RPM1 interacting protein 4; (B) *I-2*, CC-NBS-LRR, resistance protein 1; (B,C) *UDP*, *UDPI*, UDP-glucosyltransferase family 1 proteins.

**Figure 2**

**Scheme of defence responses activated by tomato (*Solanum lycopersicum*) in the presence of a complex native soil microbiota.**

(1) According to the models proposed by Chialva et al.,<sup>11</sup> in native soils microbial-associated molecular patterns (MAMPs) such as flagellin (flg22) and chitin are perceived by tomato plant. Those events elicit the PTI pathway (Plant-triggered Immunity) as a first defence level with the activation of calcium signalling (CNGCs, cyclic nucleotide-gated channels; CaM/CaM-like (CML), calmodulin-like proteins; CDPKs, calcium-dependent protein kinases) and WRKY transcription factors. This brings to the downstream activation of pathogenesis-related proteins genes (PR), such

166 as PR1, and to cell-wall fortification and lignin synthesis. (2) Since PTI-related defence is elicited, a  
 1  
 167 “continuative priming” by soil microbiota components occurs, maintaining plant defence active. (3)  
 3  
 168 When plant is attacked by *Fusarium oxysporum* f. sp *lycopersici* (FOL) the plant is already primed  
 4  
 5  
 6  
 169 and activates stronger ETI (Effector-triggered Immunity) defence. In both genotypes, effectors are  
 8  
 9  
 170 strongly perceived (e.g. by *RIN4*): only in the FOL-resistant one a specific resistance mediated by *I*-  
 10  
 11  
 171 2 is activated leading to the activation of the downstream ETI responses (such as UDP  
 13  
 14  
 172 upregulation). However, in the susceptible genotype even if *I*-2 upregulation was not observed,  
 15  
 16  
 173 FOL-suppressive soil induced the activation of downstream ETI pathway with the upregulation of a  
 18  
 19  
 174 marker UDP gene.  
 20

**Table 1. Table of primers used in RT-qPCR experiment.**


Gene	Transcript ID	Forward primer (5'-3')	Reverse primer (5'-3')	Reference
RPM1	Solyc11g0120	TCCTTCTGTAGAGTCGG	TCTTCTTCGTCGTGTTG	<sup>11</sup>
interacting protein 4 ( <i>RIN4</i> )	10.1	GCCA	GTTGGT	
CC-NBS-LRR, resistance protein 1 ( <i>I-2</i> )	Solyc11g0714	TTTGAAAGGGTCCCAA	TGCAGAGGGGTGTCAA	This study
UDP-glucosyltransferase family 1 protein (UDP)	Solyc10g0858	CAAAGCTGAAAGAGGG	TAACCCAAGCCCTAGCT	This study
UDP-glucosyltransferase family 1 protein (UDP)	80.1	AACG	CAAC	This study
UDP-glucosyltransferase family 1 protein (UDP)	Solyc09g0925	GGTGCAACCCCATGTC	ATCAGAGAATGCCGCC	This study

1	glucosyltransf	00.1	CTATTG	AAGT	study
2	erase family 1				
3	protein				
4	(UDP1)				
5					
6					
7					
8					
9					

1098

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure



Figure

